



HerbClip™

Mariann Garner-Wizard
John Neustadt, ND
Cathleen Rapp, ND

Shari Henson
Heather S Oliff, PhD
Densie Webb, PhD

Brenda Milot, ELS
Marissa Oppel, MS

Executive Editor – Mark Blumenthal

Managing Editor – Lori Glenn

Consulting Editors – Dennis Awang, PhD, Steven Foster, Roberta Lee, MD

Funding/Administration – Wayne Silverman, PhD

Production – George Solis/Kathleen Coyne

FILE: ■ Kudzu (*Pueraria montana* var. *lobata*)

■ Alcohol Consumption

■ Alcoholism

HC 070454-295

Date: December 30, 2005

RE: Kudzu Extract Found to Reduce Alcohol Consumption in Heavy Drinkers

Lukas SE, Penetar D, Berko J, et al. An extract of the Chinese herbal root kudzu reduces alcohol drinking by heavy drinkers in a naturalistic setting. *Alcohol Clin Exp Res.* May 2005;29(5):756-762.

There are no uniformly effective drugs for treating alcohol abuse or dependence. Only three drugs have been approved in the United States for treating alcoholism. Since 600 CE, herbal preparations have been used to treat alcohol-related diseases in various systems of traditional medicine. One Chinese herbal medicine, XJL (NPI-028) containing kudzu (*Pueraria montana* var. *lobata*) has been used to lessen alcohol intoxication. Kudzu is a legume with large leaves and thick long roots. Kudzu (*P. montana*) was introduced to the United States in 1876 to control soil erosion; it spread rapidly engulfing many farms. Since it has no natural predators, e.g., herbivores or insects, its spread is unchecked and the vine has become common on trees and telephone/power poles throughout the southern U.S. It has been branded "the vine that ate the South."

Previous research on Syrian golden hamsters suggested that kudzu can inhibit cravings for alcohol.¹ There are no efficacy studies on kudzu, especially as it relates to possibly reducing cravings for alcohol; most reports are anecdotal. Although the anti-intoxication mechanism of action is unclear, the isoflavones found in kudzu are believed to be responsible for the reduced alcohol intake in animals. There is one study that reported that kudzu had no effect on alcoholism in human subjects.² In addition, the authors of the present study discovered negative findings in a preliminary study. In an attempt to increase efficacy, the authors have increased the concentration of kudzu isoflavones from 1% to 25%, and the discussion below is a report of the findings. To date, other than its original intention to help stop soil erosion, there is no economic value for kudzu and it is not considered an economically important plant in the U.S. as contrasted to Asia, where its roots are commonly used in food (a thickener in sauces and soups) and as a medicinal herb.

Fourteen volunteers residing in Massachusetts with a body mass index between 19 and 24 kg/m² were screened for alcohol dependence. On the average each drank 25 alcoholic beverages each week, and none had a family history of alcoholism. The experiment was designed to determine whether pretreatment with kudzu extract alters alcohol consumption in a "natural" setting. The

natural setting was a house-like environment with a kitchen stocked with the volunteers' favorite beer and other beverages and a living room with a television, movies, etc. Volunteers had free access to the beverages and could drink at their own pace. Volunteers were given the kudzu preparation or placebo in a double-blind fashion for 7 days and then given the opportunity to drink beer in the natural laboratory. The volunteers were assessed 4 times, each separated by a washout period. One assessment was conducted before receiving kudzu and used as the baseline. Participants took 2 capsules (500 mg each) 3 times daily. Each 500 mg of kudzu (NPI-031, Natural Pharmacia Int., Inc.; Research Triangle Park, NC) contained sugar beet-based filler and 19% puerarin, 4% daidzin, and 2% daidzein. Riboflavin was added to the formulation to measure compliance. Volunteers served as their own controls.

There was a 100% compliance rate. One week treatment with kudzu extract significantly reduced the number of beers consumed, reduced volume of beer consumed, reduced the average sip size, and increased the average number of sips per beer ($P < 0.0001-0.019$). While taking kudzu, but not placebo, participants reduced their beer consumption from 6 to 4 beers per day. Data collected before each drinking session indicates that there was no change in the desire for alcohol and yet the participants drank less. There were no side effects and the patients could not identify when they were taking the active treatment. There were no biochemical changes.

The data suggests that kudzu probably does not block alcohol's effect but rather prolongs the acute effects of the first drink. This effect may reduce binge drinking. The altered sip pattern may indicate that individuals titrated (using a method or the process of determining the concentration of a dissolved substance in terms of the smallest amount of a reagent of known concentration required to bring about a given effect in reaction with a known volume of the test solution) their alcohol intake to a lower amount. The authors suggest that the first 1 or 2 beers may have satisfied their desire for beer. In the pilot study, subjects reported that kudzu plus alcohol made them feel "tired, floating, or intoxicated."

Although the actual decrease in drinking was modest, any decrease is important when the individual is drinking large quantities. The authors conclude that kudzu may help heavy or binge drinkers reduce their alcohol consumption. The lack of adverse side effects indicates that higher doses should be tested and may be more effective. This study was of the highest quality regarding design, which is not surprising considering that it was conducted by researchers from Harvard Medical School and supported by the National Institute on Alcohol Abuse and Alcoholism and the National Institute on Drug Abuse.

—Heather S. Oliff, PhD

Reference

¹Keung WM, Vallee BL. Daidzin and daidzein suppress free-choice ethanol intake by Syrian golden hamsters. *Proc Natl Acad Sci U S A*. 1993;90:10008-10012.

²Shebek J, Rindone JP. A pilot study exploring the effect of kudzu root on the drinking habits of patients with chronic alcoholism. *J Alt Compl Med*. 2000;6:45–8.

The American Botanical Council has chosen not to reprint the original article.

The American Botanical Council provides this review as an educational service. By providing this service, ABC does not warrant that the data is accurate and correct, nor does distribution of the article constitute any endorsement of the information contained or of the views of the authors.

ABC does not authorize the copying or use of the original articles. Reproduction of the reviews is allowed on a limited basis for students, colleagues, employees and/or members. Other uses and distribution require prior approval from ABC.